

Table 1. Independent determinants of homocysteine and transcobalamin in 55 hemodialysis patients

Dependent determinant	Independent determinant	Correlation coefficient	95% confidence interval	P value univariate regression	P value multiple regression ^a
Homocysteine	Creatinine	0.448	0.249, 0.685	0.0002	<0.0001
	Folate	−0.462	−0.652, −0.216	0.0005	0.0005
	Cholesterol	0.314	0.018, 0.559	0.0377	0.1268
	Triglycerides	0.279	−0.019, 0.532	0.0664	0.1417
	Albumin	0.213	−0.076, 0.469	0.1471	0.9518
	Vitamin B ₁₂	−0.167	−0.418, 0.234	0.2342	
	C-reactive prot	−0.142	−0.415, 0.155	0.3496	
	Transcobalamin	0.103	−0.184, 0.373	0.4841	
	ALAT	0.132	−0.416, 0.175	0.5562	
Transcobalamin	ALAT	0.636	0.417, 0.784	<0.0001	<0.0001
	Creatinine	0.190	−0.099, 0.450	0.1179	0.0609
	Cholesterol	−0.225	−0.486, 0.073	0.1380	0.0655
	Albumin	−0.123	−0.391, 0.164	0.4013	
	Folate	−0.117	−0.383, 0.167	0.4197	
	Homocysteine	0.103	−0.184, 0.373	0.4841	
	C-reactive prot	−0.083	−0.367, 0.216	0.5920	
	Vitamin B ₁₂	−0.076	−0.344, 0.204	0.5978	
	Triglycerides	−0.032	−0.322, 0.264	0.8375	

^aThe multiple regression analysis included only the parameters that presented a *P* value lower than 0.2 in univariate analysis.

or *MTHFR* polymorphisms, and found a weak effect of *TCN 776G* × *MTHFR 677T* allele combination on homocysteine (carriers vs. noncarriers 40.5 ± 11.3 vs. 30.9 ± 10.7 μmol/L, *P* = 0.0158). The single determinant of plasma apo-transcobalamin was alanine-aminotransferase, while creatinine was under the limit of significance (Table 1). The level was higher in ESRD, compared to 109 matched controls (719 ± 354 vs. 464 ± 160 pmol/L, *P* < 0.0001). ESRD increased plasma transcobalamin by an unknown mechanism, possibly related to glomerular filtration and tubular uptake. Under these conditions, the influence of *TCN 776C>G* on plasma transcobalamin may be limited, despite its effect on *TCN* transcription and transcobalamin intracellular concentration [3]. Therefore, in our opinion, the data of the authors are less conflicting than suggested because the weak influence of *TCN 776C>G* was initially reported in a healthy population without low serum B₁₂ [1], drug use [1], and renal dysfunction [2].

WAFAA ANWAR, PHILIPPE GÉRARD, ABDELAHIM MOUTABARREK,
FARÈS NAMOUR, and JEAN-LOUIS GUÉANT
Vandoeuvre-Nancy, France, and Casablanca, Morocco

Correspondence to Professor Jean-Louis Guéant, Laboratoire de Pathologie Cellulaire et Moléculaire en Nutrition, Equipe INSERM 00-14, Faculté de Médecine, B.P. 184, 54505 Vandoeuvre-lès-Nancy Cedex, France.
E-mail: Jean-Louis.Gueant@facmed.u-nancy.fr

REFERENCES

- WINKELMAYER WG, SKOUPY S, EBERLE C, *et al*: Effects of *TCN2 776C>G* on vitamin B₁₂, folate, and total homocysteine levels in kidney transplant patients. *Kidney Int* 65:1877–1881, 2004

- FÖDINGER M, VEITL M, SKOUPY S, *et al*: Effect of *TCN2 776C>G* on vitamin cellular availability in end-stage disease patients. *Kidney Int* 64:1095–1100, 2003
- NAMOUR F, OLIVIER JL, ABDELMOUTTALEB I, *et al*: Transcobalamin codon 259 polymorphism in HT-29 and Caco-2 cells and in Caucasians: Relation to transcobalamin and homocysteine concentration in blood. *Blood* 97:1092–1098, 2001
- ANWAR W, GUEANT JL, ABDELMOUTTALEB I, *et al*: Hyperhomocysteinemia is related to residual glomerular filtration and folates but not to methylenetetrahydrofolate-reductase and methionine synthase polymorphisms, in supplemented end-stage renal disease patients undergoing hemodialysis. *Clin Chem Lab Med* 39:747–752, 2001

Microinflammation versus inflammation in chronic renal failure patients

To the Editor: In their study, Pupim *et al* [1] show that patients with end-stage renal failure (ESRF) in the highest quartile of C-reactive protein (CRP) and interleukin-6 (IL-6) before hemodialysis (HD) initiation have a significant decrease in the serum level of both markers after 12 months of HD treatment. The baseline levels for IL-6 and CRP in these subgroups were about 58 pg/mL and 62 mg/L, respectively. As the authors note, no exclusion criteria were used for enrollment in the study. These CRP and IL-6 serum levels are probably indicative of an infection or other cause of “real inflammation” (trauma, malignancy, etc.). In the referred study microinflammation investigation is warranted. It is low-grade inflammation

that is correlated to atherosclerosis or other threatening processes. Recently, a cut-off for CRP (10 mg/L) indicative for inflammation versus microinflammation was established for the general population [2]. We also found a similar cut-off for CRP and IL-6 (9.5 mg or pg/L) indicative for an inflammatory clinical event, in a group of HD patients [3]. Possibly this cut-off is higher in other HD patients, but, as the results in the majority of studies with ESRF patients show, it seems improbable to be higher than 15 mg/L for the CRP.

The study of Pupim et al is very interesting; it shows convincingly, for the first time, that HD may not be a principal cause of inflammation and oxidative stress in ESRF patients. Although possibly the results are not essentially influenced from the above-mentioned observation, accurate assessment [4] and distinction between microinflammation and inflammation is needed in all studies in this research field.

GEORGE TSIRPANLIS, STYLIANOS CHATZIPANAGIOTOU,
and CHRYSOULA NICOLAOU
Athens, Greece

*Correspondence to George Tsirpanlis, M.D., Department of Nephrology, General Hospital of Athens, Kriezis 61, Polydrosos, Marousi, 15125, Athens, Greece.
E-mail: tsipg@hellasnet.gr*

REFERENCES

1. PUPIM LB, HIMMELFARB J, McMONAGLE E, *et al*: Influence of initiation of maintenance hemodialysis on biomarkers of inflammation and oxidative stress. *Kidney Int* 65:2371–2379, 2004
2. PEARSON TA, MENSEH GA, WAYNE AR, *et al*: Markers of inflammation and cardiovascular disease. *Circulation* 107:499–511, 2003
3. TSIRPANLIS G, BAGOS P, IOANNOU D, *et al*: Exploring inflammation in hemodialysis patients: Persistent and superimposed inflammation. A longitudinal study. *Kidney Blood Press Res* 27:63–70, 2004
4. TSIRPANLIS G, BAGOS P, IOANNOU D, *et al*: The variability and accurate assessment of the microinflammation in hemodialysis patients. *Nephrol Dial Transplant* 19:150–157, 2004